

## Report

The projects 'Impact of the trade-offs on eco-evolutionary feedbacks in predator-prey systems' and 'Indirect response to external drivers through trait variation in predator-prey systems' aimed to gain important new insights into the reciprocal interaction between trait and population dynamics in predator-prey systems. Within the projects, we quantified the distribution of several traits within green algae *Chlamydomonas reinhardtii*, the potential for trade-offs as well as the consequences of the trait variation and trade-offs for population dynamics when the green algae grow together with a rotifer predator (*Brachionus calyciflorus*) and how population dynamics contribute to the maintenance of diversity with the prey population. During the second phase of DynaTrait, we focused on the importance of the trait variation for buffering external perturbations. We used for our projects a predator-prey systems where a rotifer predator feeds on the green-algae *Chlamydomonas reinhardtii* and we focused on trait variation within *C. reinhardtii*. We used a set of previously collected *C. reinhardtii* lines from a selection experiment where different *C. reinhardtii* strains evolved with and without predation by a rotifer for 6 months. In collaboration with the groups of Ursula Gaedke (University of Potsdam) and Masato Yamamichi (University of Queensland, Australia), we combined experimental work with modelling. Together with Uwe John (AWI Bremerhaven), we complemented trait measures by sequence analyses.

### **P1.02 Becks**

#### ***Traits and trade-offs***

Shape of defence-competition trade-off: We examined the trade-off between defence and competition using isolated *Chlamydomonas reinhardtii* genotypes from a selection experiment where different *Chlamydomonas* strains (i.e., genetic background) evolved with and without predation by a rotifer for 6 months. Examining the distribution of defence against rotifer predation and maximum growth rates in trait space, we found convex trade-off curves for the isolates from the selection lines with and without rotifer predation. The trade-off curves differed, however, between the two treatments in two important ways. First, we found a significant difference in the curvature of the trade-off curve, with a stronger convex curve for the isolates for the selection lines which evolved in the presence of the rotifer predator. This more convex trade-off curve indicates the evolution of a more costly defence in the presence of the rotifer predator. Second, we found morphological differences in the evolved isolates. Isolates from the treatment evolving in the presence of the rotifer predator were growing significantly more often in aggregates of several algal cells, whereas isolates from the treatment evolving in the absence of the rotifer predator grew mostly as single cells. Additional transcriptome analyses confirmed the heritability of the phenotype. Whole genome sequencing of a subset of isolates revealed several mutations that were unique to all isolates from the selection treatment with rotifer predators but absent from the isolates from the selection treatments without rotifers. The observation of the repeated mutations suggests strong selection (Bernardes et al. 2021 *Nature Communications*).

Trait variation: In addition to defence and growth rates, we measured several additional traits to develop a better understanding of the trait variation across *C. reinhardtii* isolates. Specifically, we measured growth parameters (carrying capacity, area under the curve), morphology and the cells' ratio of carbon to nitrogen. These data haven been collected but have not been analysed yet.

Trait variation and fitness: Going one step further and aiming to link fitness with morphological and trophic traits, we measured functional response curves, half-saturation constants and morphological traits for six *C. reinhardtii* isolates. The functional response curves were used to estimate attack rates, handling time as well as pre- and post-ingestion defences (Ehrlich & Gaedke 2018 Ecol. Evol). We found that isolates can be grouped into *C. reinhardtii* with pre- and post-attack defences against consumption by the rotifer predator. Information on the morphological traits was collected by image flow cytometry (Amnis®ImageStream) and we found that size, aggregation, and circularity separated the isolates into well and less defended prey. Finally, linking the traits to fitness in the presence and absence of the rotifer predator, we found that cell size and carrying capacity explained most variation in fitness under these conditions. In addition, we used the data to test how the trade-off changes with respect to the shape of the trade-off curve (i.e., costly defence or cheap defence) and the ranking of the isolates when focusing on different traits representing defence and competitive ability. We found that the shape of the trade-off changes from between convex and linear without changing the ranking of the isolates (i.e., from most defended to least defended).

### ***Trade-offs and eco-evolutionary dynamics***

Based on theoretical work (Ehrlich et al. 2016) we designed experiments to test the invasion capability of a third algal genotype into a system with a rotifer predator and two prey genotypes. The latter are a defence or a competition specialist. The model with a two-dimensional trade-off assumes only two niches (defence and growth specialist) and the model analyse revealed that more than two types can only coexist under the assumption of fitness equality in this system. Thus, when adding a third prey genotype at low frequency to the system with the two specialists present, the possible outcomes are that the type invades and replaces one or both resident prey types or fails to invade. Whether invasion is possible or not depends on the traits of the third genotype. Initial experiments with *C. reinhardtii* and the rotifer confirmed the predictions from the model that a third type can or cannot invade depending on the trait combinations. The results of the full experiments were, however, less clear as invasion success of a third genotype differed across replicates and seemed independent of the position in trait space of the third genotype. Applying Chesson's Modern theory of Coexistence, we ran additional experimental tests with the *C. reinhardtii* isolates used in this experiment testing whether fitness and niche differences measurements were better predictors for the invasion capability than the positions in trait space. We found for most combinations of *C. reinhardtii* isolates a priority effect. As the algal and rotifer population size and the frequencies of the defended and undefended prey differed

at the time point of the addition of the third genotype, it is possible that the priority effect played an important role in this experiment. The experiments failed for technical reasons and were not further considered.

### **Development of the projects:**

Noemi Woltermann joined the project 'Impact of the trade-offs on eco-evolutionary feedbacks in predator-prey systems' as PhD student. Noemi Woltermann did however not complete her PhD thesis after becoming mother in the 2<sup>nd</sup> year of her PhD. This has slowed down the progress, but we plan to finish some of the projects with the help of Master students.

### ***P2.02 Becks***

#### ***Trait variation and buffering of external perturbations***

We tested whether and how trait variation in the prey population can buffer external perturbations to the predator using model analyses and experiments.

Indirect evolutionary rescue: Indirect evolutionary rescue (IER) is a mechanism where a non-evolving population is saved from extinction in an otherwise lethal environment by evolution in an interacting population (Yamamichi & Miner 2017). This process has been initially described for predator-prey models, where extinction of the predator is prevented by a shift in the frequency of defended towards undefended prey when reduced predator densities lower selection for defended prey. We test how increased mortality, and the initial frequencies of the prey types affect the likelihood of IER. Combining the analyses of model simulations and experiments with rotifers feeding on an algal population we show IER in the presence of increased predator mortality. We found that IER was depending on the possibility of the prey population to evolve as well as on the frequency of the defended prey, with high frequencies of defended prey resulting in extinction because prey evolution was too slow. This frequency dependency for IER was more pronounced for higher predator mortalities. (Hermann & Becks 2022 *Royal Society Open Science*).

Tracking of genotypes within mixed populations: To be able to test fundamental theories of diversity and evolutionary change often requires tracking the frequencies of clonal lineages within a population over time. We combined multispectral imaging flow cytometry and neural networks (NN) to efficiently classify clonal lineages by their heritable phenotypes and thus track their frequency changes. For this combined NN and feature values extracted from images to classify six clonal lineages of the green alga *Chlamydomonas reinhardtii* based on heritable morphological differences when paired in-silico and in-vitro. Our novel approach was able to classify clonal lineages with very high accuracy and the NN trained on pairs of clonal lineages achieved the highest accuracy of 97%. Using in-vitro samples we observed a drop in accuracy to an average of 85%, which was due to the variation occurring between image acquisitions and culturing conditions causing differences in the extracted feature values. Overall, this method can be

applied to reduce the workload and running costs of long-term lineage tracking, and it can be applied to a wide range of plankton and microbial organisms and research questions (Hermann & Becks 2024 *Methods in Ecology and Evolution*).

Contributions of ecological and evolutionary change to predator-growth rate: Evolutionary and ecological dynamics can occur on similar timescales and thus influence each other. While it has been shown that the relative contribution of ecological and evolutionary change to population dynamics can vary, it remains unknown what influences these differences. We tested whether prey populations with increased variation in their defence and competitiveness traits will have a stronger impact of evolution for predator growth rates. We controlled trait variation by pairing distinct clonal lineages of the green alga *Chlamydomonas reinhardtii* with known traits as prey with the rotifer *Brachionus calyciflorus* as predator and compared those results with a mechanistic model matching the empirical system. We measured the impact of evolution (shift in prey clonal frequency) and ecology (shift in prey population density) for predator growth rate and its dependency on trait variation using an approach based on a two-way ANOVA. Our experimental results indicated that higher trait variation, i.e., a greater distance in trait space, increased the relative contribution of prey evolution to predator growth rate over 3-4 predator generations, which was also observed in model simulations spanning longer time periods. In our model, we also observed clone-specific results, where a more competitive undefended prey resulted in a higher evolutionary contribution, independent of the trait distance. Our results suggest that trait combinations and total prey trait variation combine to influence the contribution of evolution to predator population dynamics, and that trait variation can be used to identify and better predict the role of eco-evolutionary dynamics in predator-prey systems. (Hermann, Pantel, Réveillon, Becks accepted *Journal of Evolutionary Biology*).

Microplastic and trophic interactions: While many environmental perturbations lead to the extinction of a species, some perturbations affect the trophic interactions between consumer and its resources. Microplastic affects the uptake of *C. reinhardtii* by the rotifer *B. calyciflorus* and can thus affect the trophic interaction between the two trophic levels. We tested how functional response curves for six *C. reinhardtii* isolates differing in their levels of defence and competitiveness are affected by the presence of microplastics in two different sizes (i.e., one can be ingested but not digested, the other cannot be ingested) and at different concentrations. We found that ingestion rates are affected, but that the effect was depending on the *C. reinhardtii* isolates and the microplastics sizes and concentrations. For some of the *C. reinhardtii* isolates, we found an overall reduction in the ingestion rates for all *C. reinhardtii* densities offered in the experiment, while we observed for other isolates differences only at high *C. reinhardtii* densities. The data also suggest that for some *C. reinhardtii* isolates, the ingestion rates were increased in the presence of the microplastics. In additional experiments, we found that differences in functional response curves reduced rotifer growth rates (numerical response; Réveillon & Becks 2022 bioRxiv; final revision submitted to *Ecology*).

Indirect evolutionary facilitation I: Building on the observations that the presence of microplastic affects the trophic interaction between prey and predator, we analysed a chemostat predator-prey model asking under which conditions (concentration of plastic beads and differences in the defence level of the defended prey) a shift in the prey defence level (i.e., a shift in the frequency of the defended and undefended prey) can cancel the effect the microplastics have on the predator growth rates (= indirect evolutionary facilitation (IEF)). We found that IEF occurs at high concentrations of microplastics and when the trait differences in the defence levels of the undefended and defended prey are large. For some combinations of differences in defence trait and microplastic concentrations, we observed higher predator densities in the presence of microplastics than in the absence of microplastics. The model analyses will be extended to examine how the consequences of the microplastics on the functional and numerical responses alter the dynamical behaviour of the system (e.g., shift from cycles to steady state dynamics). We performed chemostat experiments to test the predictions by the mathematical modelling (see above). The experiments confirmed the major prediction by the model, that the presence of trait variation can not only buffer the impact of an environmental disturbance (microplastics) but also it can, through the evolutionary change in the prey population, lead to higher densities in the presence of the microplastics compared to the absence of microplastics (Réveillon, Ehrlich, Gaedke, Becks *in preparation*).

Indirect evolutionary facilitation II: Expanding the work on IEF (see above), we further explored the mathematical model to test whether and when IEF can stabilize or destabilize the predator-prey dynamics, i.e., a shift from cyclic to steady state dynamics. Our analyses showed that IEF generally leads to a shift from cycles to steady state dynamics (Wonner, Réveillon, Becks, *in preparation*) (Masters thesis Tamara Wonner 2023).

#### *References*

Bernardes, J.P., John, U., Woltermann, N., Valiadi, M., Hermann, R., and L. Becks. 2021. The evolution of convex trade-offs enables the transition towards multicellularity. *Nature Communications* 12, 4222

Ehrlich, E. & U. Gaedke (2018) Not attackable or not crackable - How pre- and post-attack defences with different competition costs affect prey coexistence and population dynamics, *Ecol. Evol.* DOI: 10.1002/ece3.4145

Ehrlich, E., L. Becks, and U. Gaedke (2017) Trait-fitness relationships determine how trade-off shapes affect species coexistence. *Ecology* 98, 3188–3198

Hermann, R. J. and L. Becks. 2022. Change in prey genotype frequency rescues predator from extinction. *Royal Society Open Science*, 9, 220211.

Hermann, R. J. and L. Becks. Tracking the frequency of phytoplankton clonal lineages using multispectral image flow cytometry and neural networks. 2024. *Methods in Ecology and Evolution*, 5:401–412.

Hermann, R.J., Pantel, J.H., Réveillon, T. and L. Becks. Range of trait variation in prey determines evolutionary contributions to predator growth rates. *Journal of Evolutionary Biology* (accepted 27.4.2024)

Réveillon, T. and L. Becks. Consistency in defence and competitiveness trade-off in a planktonic predator-prey system.

bioRxiv. <https://www.biorxiv.org/content/10.1101/2022.05.02.490268v1>.

Yamamichi M, Miner BE. 2015 Indirect evolutionary rescue: prey adapts, predator avoids extinction. *Evolutionary Applications* 8, 787–795